

ICES IDENTIFICATION LEAFLETS FOR DISEASES AND PARASITES OF FISH AND SHELLFISH

No. 66

JANUARY 2017

Bonamiosis of oysters caused by *Bonamia exitiosa*

Ryan B. Carnegie



ICES
CIEM

International Council for
the Exploration of the Sea

Conseil International pour
l'Exploration de la Mer

International Council for the Exploration of the Sea Conseil International pour l'Exploration de la Mer

H. C. Andersens Boulevard 44–46
DK-1553 Copenhagen V
Denmark
Telephone (+45) 33 38 67 00
Telefax (+45) 33 93 42 15
www.ices.dk
info@ices.dk

Recommended format for purposes of citation:

Ryan B. Carnegie. 2017. Bonamiosis of oysters caused by *Bonamia exitiosa*. ICES Identification Leaflets for Diseases and Parasites of Fish and Shellfish. No. 66. 6 pp.
<http://doi.org/10.17895/ices.pub.2079>

Series Editor: Stephen Feist.

Prepared under the auspices of the ICES Working Group on Pathology and Diseases of Marine Organisms.

The material in this report may be reused for non-commercial purposes using the recommended citation. ICES may only grant usage rights of information, data, images, graphs, etc. of which it has ownership. For other third-party material cited in this report, you must contact the original copyright holder for permission. For citation of datasets or use of data to be included in other databases, please refer to the latest ICES data policy on the ICES website. All extracts must be acknowledged. For other reproduction requests please contact the General Secretary.

DOI: <http://doi.org/10.17895/ices.pub.2079>

ISBN 978-87-7482-193-9
ISSN 0109-2510

© 2017 International Council for the Exploration of the Sea

Contents

Susceptible species.....	1
Disease name.....	1
Aetiological agent.....	1
Geographical distribution.....	1
Associated environmental conditions.....	1
Significance.....	1
Gross clinical signs	2
Control measures and legislation.....	2
Key references	2
Author contact details.....	6

Bonamiosis of oysters caused by *Bonamia exitiosa*

Ryan B. Carnegie

Susceptible species

Various oysters, primarily in the genus *Ostrea* (*O. chilensis*, *O. angasi*, *O. edulis*, *O. stentina*, *O. auctorialis*, *O. equestris*, *O. puelchana*, *O. lurida*) but also *Crassostrea ariakensis* and *C. virginica* and probably *Saccostrea glomerata* (Dinamani *et al.*, 1987; Burreson *et al.*, 2004; Carnegie *et al.*, 2006; Corbeil *et al.*, 2006; Abollo *et al.*, 2008; ICES, 2013; Carnegie *et al.*, 2014; Hill *et al.*, 2014).

Disease name

Bonamiosis

Aetiological agent

Bonamia exitiosa, Phylum Haplosporidia (Carnegie *et al.*, 2000), a parasite of oyster hemocytes. Transmission is presumed to be direct.

Geographical distribution

Bonamia exitiosa was originally described from *Ostrea chilensis* in New Zealand (Dinamani *et al.*, 1987; Hine *et al.*, 2001). Since 2003, the parasite has been observed on both Atlantic and Pacific coasts of the USA, detected south of Cape Hatteras in the east (Burreson *et al.*, 2004; Dungan *et al.*, 2012) but also in Massachusetts (ICES, 2014, and in California in the west (Hill *et al.*, 2014); in southeastern Australia (Corbeil *et al.*, 2006; Carnegie *et al.*, 2014); along the Atlantic and Mediterranean coasts of Europe and North Africa, including Spain, France, the United Kingdom, Italy, Portugal and Tunisia (Abollo *et al.*, 2008; Hill *et al.*, 2010; Narcisi *et al.*, 2010; ICES, 2010; Carrasco *et al.*, 2012; Longshaw *et al.*, 2013; Batista *et al.*, 2016); and in Argentina (Kroeck and Montes, 2005).

Associated environmental conditions

Depending on the host and geographic location, clinical disease may be associated with temperatures ranging from below 10°C (Cranfield, 1968; per Hine, 1991) to over 30°C (Carnegie *et al.*, 2008). The parasite is frequently observed year-round with only a modest annual prevalence cycle displayed, as in *O. chilensis* (Hine, 1991). In *C. ariakensis*, however, prevalence and clinical disease were found to be sharply higher in the warmer summer and early fall months (Carnegie *et al.*, 2008). *B. exitiosa* displays a preference for euhaline habitats and may be inhibited by salinities below 30 ppt (Bishop *et al.*, 2006; Audemard *et al.*, 2008).

Significance

Bonamia exitiosa is acutely pathogenic in some hosts, including *O. chilensis*, *O. puelchana*, and *C. ariakensis* (Dinamani *et al.*, 1987; Burreson *et al.*, 2004; Kroeck and Montes, 2005). It can infect these species at high prevalences and intensities, causing significant mortality. From what has been reported, *B. exitiosa* is somewhat less pathogenic in other

host species. Unambiguous histological evidence of infection of *S. glomerata* from Australia and *O. auctorialis* from New Zealand is lacking although *B. exitiosa* DNA has been sequenced from both hosts (Carnegie *et al.*, 2014; Hill *et al.*, 2014) and *O. auctorialis* (as well as *O. stentina*) has been proposed to be synonymous with *O. equestris* (Shilts *et al.*, 2007), which is clearly susceptible. Effects on *C. ariakensis* and *C. virginica* have been focused primarily on young (< 1 year old) seed, with observations exclusively limited to an aquaculture context (Burrenson *et al.*, 2004; Bishop *et al.*, 2006; ICES, 2013). Because it is regarded as a significant pathogen, *B. exitiosa* looms as an impediment to fisheries and aquaculture commerce even where it is not acutely pathogenic, as for *O. edulis* in Europe and *C. virginica* in the USA. The significance of new observations is uncertain. The discovery of *B. exitiosa* in *O. edulis* in Europe, for example, may represent improved resolution of parasite diversity (*Bonamia ostreae* also being present) through the application of molecular diagnostics and DNA sequencing. The recent observation in *C. virginica* in North Carolina, USA may represent a temporary host switch under relatively stressful conditions of aquaculture, the parasite never having been observed in *C. virginica* from the region but known to infect *O. equestris* locally. Infection of *C. virginica* in Massachusetts, USA, remote from documented populations of both *B. exitiosa* and *O. equestris*, defies easy explanation.

Gross clinical signs

Bonamiosis caused by *B. exitiosa* cannot be diagnosed based on gross signs.

Control measures and legislation

Methods for control of *B. exitiosa* are not well established. Care should be taken to avoid introduction of the parasite to *B. exitiosa*-free areas. The parasite may potentially be avoided through selection of culture sites in waters of intermediate salinity (< 25, Aude-mard *et al.*, 2008) unfavourable to it, although this strategy is not practical for more stenohaline *Ostrea* species like *O. edulis*. The effectiveness and practicality of low-salinity treatment of infected oysters remains to be determined. Infection with *B. exitiosa* is a World Organisation for Animal Health (OIE)-listed disease.

Diagnostic methods


The small size of *B. exitiosa* cells (2–4 µm) makes microscopic detection challenging in cases where infection intensity is light, generally necessitating a dual strategy of detection by both microscopic and molecular means. Microscopically, *B. exitiosa* cells can be recognized in both standard histological preparations as well as stained heart or haemolymph smears as primarily uninucleate forms inhabiting the cytoplasm of oyster haemocytes and (to a lesser extent) free in oyster haemolymph. Polymerase chain reaction (PCR) assays specific for *B. exitiosa* have been developed (Carnegie *et al.*, 2008; Ramilo *et al.*, 2013), the latter adaptable for use in a SYBR Green real-time PCR format or multiplexed in conventional PCR format with an assay presented in the same publication for *B. ostreae*. An older conventional assay detecting *B. exitiosa* via PCR-restriction fragment length polymorphism (RFLP, Hine *et al.*, 2001) remains in wide use. Available *in situ* hybridisation (ISH) assays (Cochennec *et al.*, 2000; Carnegie *et al.*, 2003) remain genus-specific at best. Transmission electron microscopy (TEM) is not a practical tool for *B. exitiosa* diagnosis.

Key references


Abollo, E., Ramilo, A., Casas, S. M., Comesaña, P., Cao, A., Carballal, M. J., Villalba, A. 2008. First detection of the protozoan parasite *Bonamia exitiosa* (Haplosporidia) infecting flat oyster *Ostrea edulis* grown in European waters. *Aquaculture*, 274: 201-207.

- Audemard, C., Carnegie, R. B., Bishop, M. J., Peterson, C. H., Bureson, E. M. 2008. Interacting effects of temperature and salinity on *Bonamia* sp. parasitism in the Asian oyster *Crassostrea ariakensis*. *Journal of Invertebrate Pathology*, 98: 344-350.
- Bishop, M. J., Carnegie, R. B., Stokes, N. A., Peterson, C. H., Bureson, E. M. 2006. Predicting the outcome of a non-native oyster introduction: facilitation of an enzootic parasite. *Marine Ecology Progress Series*, 325: 145-152.
- Batista, F. M., López-Sanmartín, M., Grade, A., Navas, J. I., Ruana, F. 2016. Detection of *Bonamia exitiosa* in the European flat oyster *Ostrea edulis* in southern Portugal. *Journal of Fish Diseases*, 39: 607-611.
- Bureson, E. M., Stokes, N. A., Carnegie, R. B., Bishop, M. J. 2004. *Bonamia* sp. (Haplosporidia) found in non-native oysters, *Crassostrea ariakensis*, in Bogue Sound, North Carolina. *Journal of Aquatic Animal Health*, 16: 1-9.
- Carnegie, R. B., Barber, B. J., Distel, D. L. 2003. Detection of the oyster parasite *Bonamia ostreae* by fluorescent *in situ* hybridization. *Diseases of Aquatic Organisms*, 55: 247-252.
- Carnegie, R. B., Hill, K. M., Stokes, N. A., Bureson, E. M. 2014. The haplosporidian *Bonamia exitiosa* is present in Australia, but the identity of the parasite described as *Bonamia roughleyi* is uncertain. *Journal of Invertebrate Pathology*, 115: 33-40.
- Carnegie, R. B., Barber, B. J., Culloty, S. C., Figueras, A. J., Distel, D. L. 2000. Development of a PCR assay for detection of the oyster pathogen *Bonamia ostreae* (Pichot *et al.*, 1980), and support for its inclusion in the Haplosporidia. *Diseases of Aquatic Organisms*, 42: 199-206.
- Carnegie, R. B., Bureson, E. M., Hine, P. M., Stokes, N. A., Audemard, C., Bishop, M. J., Peterson, C. H. 2006. *Bonamia perspora* n. sp. (Haplosporidia), a parasite of the oyster *Ostreola equestris*, is the first *Bonamia* species known to produce spores. *Journal of Eukaryotic Microbiology*, 53: 232-245.
- Carnegie, R.B., Stokes, N.A., Audemard, C., Bishop, M.J., Wilbur, A.E., Alphin, T.D., Posey, M.H., *et al.* 2008. Strong seasonality of *Bonamia* sp. infection and induced *Crassostrea ariakensis* mortality in Bogue and Masonboro Sounds, North Carolina, USA. *Journal of Invertebrate Pathology*, 98: 335-343.
- Carrasco, N., Villalba, A., Andree, K. B., Engelsma, M. Y., Lacuesta, B., Ramilo, A., Gairín, I., *et al.* 2012. *Bonamia exitiosa* (Haplosporidia) observed infecting the European flat oyster *Ostrea edulis* cultured on the Spanish Mediterranean coast. *Journal of Invertebrate Pathology*, 110: 307-313.
- Cochennec, N., Le Roux, F., Berthe, F., Gérard, A. 2000. Detection of *Bonamia ostreae* based on small subunit ribosomal probe. *Journal of Invertebrate Pathology*, 76: 26-32.
- Corbeil, S., Arzul, I., Robert, M., Berthe, F. C. J., Besnard-Cochennec, N., Crane, M. St. J. 2006. Molecular characterization of an Australia isolate of *Bonamia exitiosa*. *Diseases of Aquatic Organisms*, 71: 81-85.
- Cranfield, H. J. 1968. An unexploited population of oysters, *Ostrea lutaria* Hutton, from Foveaux Strait. *New Zealand Journal of Marine and Freshwater Research*, 2: 3-22.
- Dinamani, P., Hine, P. M., Jones, J. B., 1987. Occurrence and characteristics of the haemocyte parasite *Bonamia* sp. in the New Zealand dredge oyster *Tiostrea lutaria*. *Diseases of Aquatic Organisms*, 3: 37-44.
- Dungan, C. F., Carnegie, R. B., Hill, K. M., McCollough, C. B., Laramore, S. E., Kelly, C. J., Stokes, *et al.* J. 2012. Diseases of *Crassostrea ariakensis* and *C. virginica* oysters reared in ambient waters from the Choptank River, Maryland and the Indian River Lagoon, Florida. *Diseases of Aquatic Organisms*, 101: 173-183.
- Hill, K. M., Carnegie, R. B., Aloui-Bejaoui, N., El Gharsalli, R., White, D., Stokes, N. A., Bureson, E. M. 2010. Observation of a *Bonamia* sp. infecting the oyster *Ostrea stentina* in Tunisia, and a consideration of its phylogenetic affinities. *Journal of Invertebrate Pathology*, 103: 179-185.

- Hill, K. M., Stokes, N. A., Webb, S. C., Hine, P. M., Kroeck, M. A., Moore, J. D., Morley, M. S., *et al.* 2014. Phylogenetics of *Bonamia* parasites based on small subunit and internal transcribed spacer region ribosomal DNA sequence data. *Diseases of Aquatic Organisms*, 110: 33-54.
- Hine, P. M. 1991. The annual infection pattern of infection by *Bonamia* sp. in New Zealand flat oysters, *Tiostrea chilensis*. *Aquaculture*, 93: 241-251.
- Hine, P. M., Cochenne-Laureau, N., Berthe, F. C. J. 2001. *Bonamia exitiosus* n. sp. (Haplosporidia) infecting flat oysters *Ostrea chilensis* in New Zealand. *Diseases of Aquatic Organisms*, 47: 63-72.
- ICES. 2010. Report of the Working Group on Pathology and Diseases of Marine Organisms (WGPDMO), 23-27 February 2010, Uppsala, Sweden. ICES CM 2010/SSGHIE:02. 66 pp.
- ICES. 2013. Report of the Working Group on Pathology and Diseases of Marine Organisms (WGPDMO), 5-9 March 2013, Padova, Italy. ICES CM 2013/SSGHIE:03. 30 pp.
- ICES. 2014. Second interim report of the Working Group on Pathology and Diseases of Marine Organisms (WGPDMO), 25-28 February 2014, ICES Headquarters, Copenhagen, Denmark. ICES CM 2014/SSGHIE:02. 26 pp.
- Kroeck, M. A., Montes, J. 2005. Occurrence of the haemocyte parasite *Bonamia* sp. in flat oysters *Ostrea puelchana* farmed in San Antonio Bay (Argentina). *Diseases of Aquatic Organisms*, 63: 231-235.
- Longshaw, M., Stone, D. M., Wood, G., Green, M. J., White, P. 2013. Detection of *Bonamia exitiosa* (Haplosporidia) in European flat oysters *Ostrea edulis* cultivated in mainland Britain. *Diseases of Aquatic Organisms*, 106: 173-179.
- Narcisi, V., Arzul, I., Cargini, D., Calzetta, A., Traversa, D., Robert, M., Joly, J. P., *al.* 2010. Detection of *Bonamia ostreae* and *B. exitiosa* (Haplosporidia) in *Ostrea edulis* from the Adriatic Sea (Italy). *Diseases of Aquatic Organisms*, 89: 79-85.
- Ramilo, A., Navas, J. I., Villalba, A., Abollo, E. 2013. Species-specific diagnostic assays for *Bonamia ostreae* and *B. exitiosa* in European flat oyster *Ostrea edulis*: conventional, real-time and multiplex PCR. *Diseases of Aquatic Organisms*, 104: 149-161.
- Shilts, M. H., Pascual, M. S., Ó Foighil, D. 2007. Systematic, taxonomic and biogeographic relationships of Argentine flat oysters. *Molecular Phylogenetics and Evolution*, 44: 467-473.



Histological section of phloxine-tartrazine-stained tissue from a *Bonamia exitiosa*-infected oyster (*Crassostrea ariakensis*) showing parasite cells both intrahaemocytic and free in haemal spaces of connective tissues. Bar = 10 microns. (Photo: R. B. Carnegie, VA Institute of Marine Science. Histology courtesy of C. F. Dungan, Maryland Department of Marine Resources, USA).



Cytological preparation of Hemacolor (Merck)-stained oyster hemolymph from a *Bonamia exitiosa*-infected oyster (*Crassostrea ariakensis*) showing parasite cells distributed extracellularly among oyster haemocytes. Bar = 10 microns. (Photo: R. B. Carnegie, VA Institute of Marine Science).

Author contact details

Ryan B. Carnegie

Virginia Institute of Marine Science

College of William & Mary

P.O. Box 1346

Gloucester Point, Virginia 23062

USA

carnegie@vims.edu